

File Specification for GEOS-DAS Gridded Output Version 5.3

*Global Modeling and Assimilation Office
Goddard Space Flight Center, Greenbelt, Maryland*

<http://gmao.gsfc.nasa.gov>

Release Date March 30, 2004; Expiration Date March 30, 2009

Version 5.3
To support EOS with GEOS-4.x

Document maintained by Rob Lucchesi (GMAO, SAIC)
Permission to quote from this document should be obtained from the GMAO.



**Michele Rienecker
Global Modeling and Assimilation Office
Earth Sciences Directorate
NASA Goddard Space Flight Center
Greenbelt, Maryland 20771**

This page intentionally left blank.

File Specification for GEOS-DAS Gridded Output

Document No. GMAO-1001v5.3

March 30, 2004

Approved by:

Michele Rienecker Date
Head
Global Modeling and Assimilation Office
Code 900.3, NASA GSFC

Gi-Kong Kim Date
CCB Chair
Global Modeling and Assimilation Office
Code 902.0, NASA GSFC

REVISION HISTORY

Version Number	Revision Date	Pages Affected/Extent of Changes	Approval Authority
Version 1.0 (original - no electronic copy)	April 8, 1997	All	Not Approved
Version 3.0 (DAO-1001v3.0)	November 4, 1997	<ul style="list-style-type: none"> - Added 5 authors - Reorganized the document - Replaced Sections 3, 5, 6, and Appendix B 	Not Approved
Version 3.1 (DAO-1001v3.1)	January 5, 1998	<ul style="list-style-type: none"> - Added Figure 1 - Revised size estimate for single level mis and lsm in Sections 6.1 and 6.2 - Made various changes to the list of variables in the assimilated time averaged files in Section 6.3 	Rob Lucchesi 1/26/99 (Closed CR 131)
Version 3.2 (DAO-1001v3.2)	March 4, 1999	<ul style="list-style-type: none"> - Changed "GEOS-3" to "GEOS-AM1" throughout the document - Updated scope and launch date in Section 1 - Provided more detailed description of file format including Dimensions, Variables, and Attributes in Section 2 - Modified the horizontal resolution and clarified the description of the grids in Section 4 - Included version ID in the filename convention described in Section 5.1 - The file sizes were updated in Section 6 to reflect the new horizontal resolution - Improved the description of packing and included directions for unpacking in Section 8 - Two sample programs were added in Section 9 to demonstrate using the HDF library and the HDF-EOS library for reading the data files - Updated the description of First Look Analysis and added a description of Late Look Analysis in Appendix A - Added references for the data format 	CCB 3/16/99
Version 4.0 (DAO-1001v4.0)	May 19, 1999	<ul style="list-style-type: none"> - Title changed to "File Specification for GEOS-DAS Gridded Output" - Changed details of internal products in Section 1 - Added a figure and description of the horizontal grid structure, removed references to sigma levels in the vertical grid description, and indicated differences in the grids of the CHM products in Section 4 - Indicated that forecast and analysis products will not be archived for distribution by ECS and removed file-naming information for sigma and forecast products in Section 5 - Various changes made to Section 6: - split product table into one table for first look and one for late look 	CCB 5/19/99

Version Number	Revision Date	Pages Affected/Extent of Changes	Approval Authority
		<ul style="list-style-type: none"> - removed sigma products and sigma sizes and definitions - added ESDTs - adjusted the horizontal resolution of CHM products - removed forecast products - added a discussion of the different horizontal and vertical grid structures of the CHM products - Updated the description of First Look Analysis in Appendix A - Added a new table of pressure levels for CHM products and removed the table of sigma levels in Appendix C - Added references for the data format - Levels 36 and 42 in Appendix C tables were changed from 0.1 to 0.2 because in 3D pressure products, 0.2 hPa is now the highest level, changed from 0.1 hPa in earlier documentation 	
Version 4.1 (DAO-1001v4.1)	August 18, 1999	- Slight adjustment of variable content for tavg_cld_p, tavg3d tmp_p, and tavg3d tmp_p products.	CCB 8/18/99
Version 4.2 (DAO-1001v4.2)	January 21, 2000	<ul style="list-style-type: none"> - Re-write Section 3 to clarify description of time-averaged products. - Changed AM-1 to Terra throughout document. - Added approval dates to revision history for Version 4.0 and Version 4.1. 	CCB 1/21/00
Version 4.3 (DAO-1001v4.3)	September 20, 2001	<ul style="list-style-type: none"> - In Table 6.1, adjusted the sizes of three products to match the detailed descriptions in Section 6.2: DFLAXENG changes from 66.7 to 58.4 DLLAPCLD changes from 150.1 to 112.6 DLLAPTMP changes from 337.8 to 412.9 	CCB 5/17/00
Version 5.0 <i>DRAFT</i> in support of GEOS-4.x.	November 07, 2001	<ul style="list-style-type: none"> -The horizontal resolution of output products has been changed to 288x181. The prior value was 360x181. This change is reflected in Sections 4 and 6. -The grid origin has been changed from (-180.0W,-90.0S) to (0.0W,-90.0S). This change is documented in Section 4. -Adjustments were made to Section 3 to describe the different time-averaging characteristics of the GEOS-4.x system output variables. In particular, 6-hourly time averages now have a different averaging period. This will result in different range begin and end times for 3D pressure products. -In section 6, variables that were unavailable from the GEOS-4.x system were removed from DAS product descriptions. Variables that were previously produced by GEOS-3.x but are no longer available are indicated with a strike-through. Similarly, the old file size calculations are indicated with strike-throughs. -Appendix A was cleaned up to better describe the assimilation modes that are used in the operational arena. -Figures specific to GEOS-3.x were removed. 	DDB 10/31/01
Version 5.0 in support	March 20, 2002	-Time-averaging definitions were changed. All time-	CCB 3/22/02

Version Number	Revision Date	Pages Affected/Extent of Changes	Approval Authority
of GEOS-4.x (DAO-1001v5.0)		<p>averages will be time-stamped with the center of the averaging period. The range time for each time-averaged product will be 00GMT to 00 GMT. This is different from GEOS-3, where products had range times from 21 GMT to 21 GMT. Users will have adjust indices used to read time in order to get data valid for the same time period.</p> <ul style="list-style-type: none"> -Z0 replaced by Z0H and Z0M in tavg2d_str_x. -GWET in tsyn2d_mis_x was replaced by GWETROOT and GWETTOP in tavg2d_eng_x. -Variable name TGROUND changed to TSKIN in tsyn2d_mis_x and tavg2d_eng_x. -OMEGA moved from tsyn3d_mis_p to tavg3d_mom_p. -Added PRECL to tavg2d_eng_x. PREACC = PRECL + PRECON. -Moved PARDR and PARDF from tavg2d_lsm_x to tavg2d_eng_x. -Removed from tsyn2d_mis_x: ALBEDO, VAVEU, VAVEV, VAVET, TPW, GWET, SNOW -Removed from tsyn3d_mis_p: TKE, OMEGA. -Removed from tavg2d_eng_x: QICE, CT, VAVEQIAU, VAVEQFIL, VAVETIAU. -Removed from tavg2d_str_x: GWDUT, GWDVT, CU, Z0, PIAU. -Removed from tavg2d_lsm_x: SNOWFALL, GREEN, DLWDTC, PARDR, PARDF. -Removed from tavg2d_cld_x: TAULOW, TAUMID, TAUHI, CLDTMP, CLDPRS. -Removed from tavg3d_cld_p: CLDRAS. -Removed from tavg3d_mom_p: RFU, RFV, UIAU, VIAU, DUDT, DVDT. -Removed from tavg3d_mst_p: DQLS, QIAU, QFILL, DQDT. -Removed from tavg3d_tmp_p: TURBT, MOISTT, RADLW, RADSW, GWDT -Added TROPQ, UFLUX, VFLUX, and HFLUX to tsyn2d_mis_x; -PBL replaced by PBLH. 	
Version 5.1 (DAO-1001v5.1)	April 25,2002	The longitude range of the grid was changed from (0 -> 360) to (-180 -> 180). This is consistent with the GEOS-3 products and with ECS conventions.	DDB 04/25/2002
Version 5.2 (DAO-1001v5.2)	May 23, 2002	<ul style="list-style-type: none"> -The filename examples in section 5 were updated to reflect the new range times of time-averaged products. -Appendix D was added highlighting the differences between GEOS-3 and GEOS-4 products. 	CCB off-line 5/23/2002
Version 5.3 (GMAO-1001v5.3)	February 19, 2004	<ul style="list-style-type: none"> - Appendix E added to describe two new ETA-coordinate products. Section 5 modified to add ETA terminology to file and ESDT naming convention. - Data Assimilation Office references changed to Global 	CCB, off-line 3/30/2004

Version Number	Revision Date	Pages Affected/Extent of Changes	Approval Authority
		Modeling and Assimilation Office. - Typo in Section 3 (01:30 changed to 13:30). - Reformatted Appendix D for clarity and included PARDR and PARDF.	

Table of Contents

1. Introduction.....	3
2. Format and File Organization	3
<i>2.1 Dimensions.....</i>	<i>4</i>
<i>2.2 Variables.....</i>	<i>4</i>
<i>2.3 Global Attributes.....</i>	<i>6</i>
3. Synoptic vs Time-averaged Products.....	7
4. Grid Structure.....	8
5. File Naming Convention.....	9
<i>5.1 File Name.....</i>	<i>9</i>
<i>5.2 Earth Science Data Types (ESDT) Name</i>	<i>11</i>
6. File Collections	13
<i>6.1 File Sizes</i>	<i>13</i>
<i>6.2 Assimilated Synoptic Files</i>	<i>13</i>
<i>6.3 Assimilated Time Averaged Files.....</i>	<i>16</i>
<i>6.4 Assimilated Chemistry Files</i>	<i>20</i>
7. Metadata	21
<i>7.1 EOSDIS Metadata.....</i>	<i>21</i>
<i>7.2 COARDS Metadata.....</i>	<i>21</i>
8. Sample Software.....	23
Appendix A. Types of Assimilation Configurations	29

Appendix B. Collection Metadata 30

Appendix C. Vertical Grid Structure..... 34

Appendix D. Summary of differences between GEOS-3 and GEOS-4..... 35

Appendix E: Additional products on eta vertical coordinates..... 38

REFERENCES..... 42

1. Introduction

This document describes the gridded output files from the version of the Goddard EOS Data Assimilation System (GEOS-4), which will support level-4 product generation. The intended audience is EOS instrument teams and other users of GEOS-4 products who need to write software to read GMAO products. The gridded data described in this document will be produced by the GEOS-4 Data Assimilation System (DAS) beginning in 2002 and delivered to the Goddard Distributed Active Archive Center (DAAC), part of the EOSDIS Core System (ECS). Two production suites will run daily, the first-look assimilation and the late-look assimilation (see Appendix A for definitions). To conserve space, a limited set of products will be generated by the first-look assimilation while the full set will be generated by the late-look assimilation. The product descriptions in Section 6 will identify which products are generated by each run. Information on the status of GMAO product generation can be found at <http://gmao.gsfc.nasa.gov/Operations/>.

2. Format and File Organization

GEOS- 4 files are in HDF-EOS format, which is an extension of the Hierarchical Data Format (HDF), Version 4 developed at the National Center for Supercomputing Applications (NCSA). Each GEOS- 4 file will contain a single HDF-EOS grid, which in turn contains a number of geophysical quantities that we will refer to as "fields" or "variables." Some files will contain 2-D variables on a lon/lat grid and some files will contain 3-D variables on the same lon/lat grid but with an additional vertical dimension. All files will have a time dimension with the number of times dependent on the file type (see Section 6). GEOS- 4 products always contain one complete day of data.

The variables are created using the **GDdefield** function from the HDF-EOS GD API which implements them as HDF Scientific Data Set (SDS) arrays so they can be read with standard HDF routines. In addition to the geophysical variables, the files will have SDS arrays that define dimension scales (or coordinate variables). There will be two distinct scales for each dimension, which will insure that a wide variety of graphical display tools can interpret the dimension scales. In particular, there is a set of dimension scales that adhere to the COARDS conventions (see References).

ECS metadata and other information will be stored as global attributes.

2.1 Dimensions

GMAO HDF-EOS files will contain two sets of dimension scale (coordinate) information. One set of dimensions is defined using the **SDsetdimscale** function of the standard HDF SD interface. This set of scales will have an attribute named "units", set to an appropriate string defined by the COARDS conventions that can be used by applications to identify the dimension. The other set of dimension scales is created using the **GDddefield/GDwritefield** functions as suggested in the ECS technical paper "Writing HDF-EOS Grid Products for Optimum Subsetting Services."

Table 2.1-1. Dimension Variables Contained in GMAO HDF-EOS Files

Name	Description	Type	units attribute
XDim:EOSGRID	longitude values	float32	degrees_east
YDim:EOSGRID	latitude values	float32	degrees_north
Height:EOSGRID (3D only)	pressure levels	float32	millibar
TIME:EOSGRID	minutes since first time in file	float32	minutes since YYYY-MM-DD HH:MM:SS
XDim	longitude values	float64	N/A
YDim	latitude values	float64	N/A
Height (3D only)	pressure levels	float64	N/A
Time	seconds since 1/1/93	float64	N/A

The 32-bit dimension variables have a "units" attribute that makes them COARDS-compliant, while the 64-bit dimension variables satisfy ECS requirements.

2.2 Variables

Variables are stored as SDS arrays even though they are defined with the HDF-EOS **GDddefield** function. As a result, one can use the SD interface of the HDF library to read any variable from the file. The only thing one must know is the short name of the variable and the dimensions. You can quickly list the variables in the file by using common utilities such as *ncdump* or *hdp*. The latter utility is distributed from NCSA with the HDF library. A sample scan of one GMAO HDF-EOS file is shown below:

```
unix% hdp dumpsds -h
DAS.fk.asm.tsyn3d_mis_p.GEOS400.1999052800.1999052818.V01 | grep Variable
```

```
Variable Name = UWND
Variable Name = VWND
Variable Name = HGHT
Variable Name = TMPU
Variable Name = SPHU
Variable Name = RH
Dimension Variable Name = XDim:EOSGRID
Dimension Variable Name = YDim:EOSGRID
```

Dimension Variable Name = Height:EOSGRID
 Dimension Variable Name = TIME:EOSGRID
 Variable Name = XDim
 Variable Name = YDim
 Variable Name = Height
 Variable Name = Time

In Section 8 we will present sample code for reading one or more data fields from this file. The short names for all variables in all GMAO data products are listed in the File Collections chapter, Section 6.

Each variable will have metadata attributes defined that may be useful. Many of these attributes are required by the COARDS conventions while others are for internal GMAO use. A listing of required attributes follows:

- FillValue (Type = 32-bit floating point or 16-bit signed integer)
 Any missing data will be filled with this value. This attribute is generated automatically by the HDF-EOS library. It will be of the same data type as the variable, thus could be 16-bit integer if the variable is packed.
- long_name (Type = 8-bit signed char)
 Long name or description for this variable.
- units (Type = 8-bit signed char)
 The units of the geophysical variable (ud-units compliant).
- scale_factor (Type = 32-bit floating point)
 If this variable is packed into 16-bit integers, this value would be multiplied to each data value to obtain the floating point number.
- add_offset (Type = 32-bit floating point)
 For a packed variable, add this number after multiplying by the scale_factor.
- missing_value (Type = 32-bit floating point or 16-bit signed integer)
 For an unpacked variable, this is a floating-point representation of the fill value. For a packed variable, this is an integer representation of the fill value.
- fmissing_value (Type = 32-bit floating point)
 Always a floating-point representation of the fill value.
- vmin (Type = 32-bit floating point)
 For internal GMAO use.
- vmax (Type = 32-bit floating point)
 For internal GMAO use.

Since we are not planning to pack the products intended for use by EOS instrument teams, attributes that support packing can be ignored. There may be other attributes that are meant for internal GMAO use that can also be ignored.

2.3 Global Attributes

In addition to SDS arrays containing variables and dimension scales, there is additional metadata stored in GMAO HDF-EOS files. Some of this metadata is required by the COARDS conventions, some is present due to ECS requirements, and some may exist as a convenience to internal GMAO users. A summary of global attributes that will exist in all GMAO files follows:


- HDFEOSVersion - Version of the HDF-EOS library used to create this file.
- StructMetadata.0 - This is the GridStructure metadata that is created by the HDF-EOS library.
- CoreMetadata.0 - The ECS inventory metadata.
- ArchivedMetadata.0 - The ECS archive metadata.
- Title - GEOS-DAS Version number and CVS tag.
- Conventions - Should always be set to "COARDS".
- Source - NASA Global Modeling and Assimilation Office, GEOS-DAS Version.
- Contact - Where to send questions.
- History - Version of DAS output library.

3. Synoptic vs Time-averaged Products

GEOS-4 gridded output files are identified as either synoptic or time-averaged products. Synoptic products are defined in Section 6.2 while time-averaged products are defined in Section 6.3. Additional synoptic products associated with the ozone assimilation are defined in Section 6.4. All synoptic products contain fields that are snapshots of a specific time. The synoptic products defined in Section 6.2 are generated directly from the atmospheric data analysis, which is performed every six hours at the four *synoptic times* (00 GMT, 06 GMT, 12 GMT, and 18 GMT). The "tsyn3d_mis_p" product has a time frequency of 6 hours, with data for each synoptic time. The "tsyn2d_mis_x" synoptic product has a time-frequency of 3 hours. The values for 03 GMT, 09 GMT, 15 GMT, and 21 GMT are time-interpolated from the synoptic times. This is done to make the product consistent with GEOS-3 products.

Time-averaged products are averaged over a 3-hour period for single level files or a 6-hour period for pressure level files. Single level products contain 8 times, with time-stamps of 01:30, 04:30, 07:30, 10:30, 13:30, 16:30, 19:30, and 22:30 GMT. Each time-stamp represents the center of a three-hour average, i.e., the first time in each file (time stamped with 01:30 GMT) represents the average between 00 GMT and 03 GMT. Time-averaged pressure level products contain 4 times with time-stamps of 03, 09, 15, and 21 GMT. Each time-stamp represents the center of a 6-hour average, i.e., the first time in the file (time-stamped with 03 GMT) represents an average between 00 GMT and 06 GMT. The valid range time of a daily file is 00 GMT to 00 GMT for both 3-hour and 6-hour average products. Note that the time-averaging of GEOS-4 products is different from the preceding GEOS-3 system (see diagram). Also note that eta-level products, added to the GEOS-4 product suite in 2004, have a different time-averaging convention. Please see Appendix E for details.

DAS Time-averaged Products



4. Grid Structure

Most GEOS-4 gridded output will be on a 1.25 x 1.0 degree global longitude-latitude horizontal grid (288,181). Products from the GMAO ozone assimilation (D4FAXCHM, D4FAPCHM, D4LAXCHM, D4LAPCHM) will be on a 2.5 x 2.0 degree longitude-latitude horizontal grid (144,91). The horizontal grid origin is the lower-left point and represents the geolocation (-180.0 W, -90.0 S). Latitude or longitude as a function of the index can be determined by:

$$\text{lon} = -180 + (x-1) * dx$$

$$\text{lat} = -90 + (y-1) * dy$$

where $dx = 1.25$, $dy = 1.0$ except in the case of ozone products where $dx = 2.5$, $dy = 2.0$. For all parameters of each file, the gridpoints represent the center of a box, i.e., the value at (lon=5,lat=0) represents a box bounded by the points (lon=4.375,lat=-0.5), (lon=5.625,lat=0.5), (lon=5.625,lat=0.5), and (lon=4.375,lat=-0.5). Scalar values are the volume mean within the box.

The vertical structure of gridded products will have 3 different configurations: single-level (can be vertical averages), pressure levels, or eta levels (discussed in Appendix E). Single level data for a given variable appear as 3-dimensional fields (x,y,time) while pressure level data appear as 4-dimensional fields (x,y,z,time). Pressure level data will be output on 36 pressure levels (hPa), except for D4FAPCHM and D4LAPCHM which will have 42 pressure levels (see Appendix C). The appropriate grid structure will be specified both in the filename and the metadata.




Figure 4-1: 1.25 x 1.0 degree horizontal grid.
 The origin is at lower-left corner (-180W, -90S).
 $\text{lon} = -180 + (x-1) * dx$
 $\text{lat} = -90 + (y-1) * dy$

5. File Naming Convention

Each GEOS-4 product will have a complete file name identified in the EOSDIS metadata as "LocalGranuleID". EOSDIS also requires abbreviated naming indices (8-character limit) for each Earth Science Data Type (ESDT). The ESDT indices convention is described in section 5.2.

5.1 File Name

The standard generic complete name for the assimilated GEOS- 4 configuration products will appear as follows:

DAS.config.mode.filetype.expid.yyyymmddhh.yyyymmddhh.version

A brief description of the node fields appear below:

DAS:

Identifies output as a Data Assimilation System product.

config:

GEOS-4 will run in two different configurations. Appendix A describes these configurations. These are the strings that can appear in the config field.

flk - First Look

llk - Late Look

mode:

GEOS-4 can run in different modes of operation, but the only mode used to support EOS instrument teams is "asm".

asm - Assimilation. Uses a combination of atmospheric data analysis and model forecasting to generate a time-series of global atmospheric quantities.

filetype:

The major filetypes are subdivided into file collections. Collections contain several fields with common characteristics. These collections are necessary to keep file sizes reasonable. Each filetype will contain the following information:

type/dimension_group_level

type/dimension:

There exist four possible type/dimension conventions for the DAS data products:

tsyn2d - 2-dimensional instantaneous fields at synoptic times (no time averaging).

tsyn3d - 3-dimensional instantaneous fields at synoptic times (no time averaging).

tavg2d - 2-dimensional 3-hour time averaged fields, time-stamped at the center of the averaging period. For example, 04:30z output would be a 3z-6z time average).

tavg3d - 3-dimensional 6-hour time averaged fields, time-stamped at the center of the averaging period. For example, 3z output would be a 0z-6z time average.

group (for files which contain 2-dimensional fields):

str: stress related fields

eng: energy related fields

cld: cloud related fields

lsm: land surface model related fields

chm: chemical related fields (e.g. ozone)

mis: mixture of prognostic and diagnostic fields

group (for files which contain 3-dimensional fields):

mis: mixture of prognostic and diagnostic fields

mom: momentum profile fields

tmp: temperature profile fields

mst: moisture profile fields

cld: cloud profile related fields

trp: transport profile fields (e.g. eddy diffusivity)

chm: chemical profile fields

level: There are three possible level types for the DAS data:

x: single level data (surface, column integrated, single level)

p: pressure level data (see Appendix C for pressure levels)

e: ETA level data (see Appendix E)

expid:

Experiment Identification. The GEOS-4 DAS data sets will be labeled:

GEOS4##

where ## is a two-digit number. At the time of this document's release, the experiment identification is GEOS403. When GEOS-4 was initialized in operations, the experiment identification was GEOS401. When a reprocessing experiment is run over previously processed data periods or when a patched version is put into First-look or Late-look operations, we will increment the ## sequentially. Note that products produced by GEOS-3 had an experiment identification of AM101. These First-look and Late-look products were generated between January 1, 2000 and the operations date of GEOS-4 (October 1, 2002). As updated versions of the GEOS software are implemented in operations, the cvs tag in the metadata parameter "Title" will be

modified as well as the GMAO operations status web page (<http://gmao.gsfc.nasa.gov/Operations/>).

yyyymmddhh.yyyyymmddhh:

This group defines the beginning date and synoptic time as well as the ending date and synoptic time.

yyyy - year string (e.g. "2002")

mm - month string (e.g. "09" for September)

dd - day of the month string (e.g. "10" for the tenth day of the month)

hh - time (synoptic hour, e.g. "18" for 18:00 Greenwich Mean Time)

version:

This group defines the file version and takes the form V##. Under normal conditions ## will be 01. In the event of a processing error that requires a re-processing, this number will be incremented to identify the new version of this file. The file version will also be represented in the EOSDIS metadata as "LocalVersionID".

EXAMPLE:

DAS.flk.asm.tavg3d_mom_p.GEOS401.2002091500.2002091600.V01

This is an example of a DAS filename. It is a first look assimilation product. The data are 6-hour time averaged output on pressure levels (3 dimensions). The filetype consists of momentum fields. The valid range time for the file is Sep 15 at 00 GMT through Sep 16 at 00 GMT. There are 4 times, each representing a 6-hour average. The first average is from 00 GMT to 06 GMT. The last average is from 18 GMT to 00 GMT. See the discussion on time-averaged data in section 3 for more information.

5.2 Earth Science Data Types (ESDT) Name

To accommodate EOSDIS toolkit requirements, GEOS-4 complete filenames are associated to shorter or abbreviated indices in the ESDTs. EOSDIS requires a short (8 character) name for each ESDT. Below is the abbreviated naming convention for the GEOS-4 gridded ESDTs. The standard ESDT naming convention for the GEOS-4 gridded output will have the form:

DSPMVCCC

D: DAS identifier. Always **D**.

S: Major system number. For GEOS-4, **4**.

P: Product

F = First Look

L = Late Look

M: Mode

A = Assimilation

F = Forecast (currently not used)

V: Vertical Coordinate:

X = Single-Level

P = Pressure

E = ETA (Appendix E)

CCC: Filetype

MIS = misc

ENG = energy

LSM = land surface model

STR = stress

CHM = chemical

CLD = cloud

MOM = momentum

MST = moisture

TMP = temperature

TRP = transport fields

Example:

Abbreviated Name: D4FAPMOM

Complete Name: DAS.flk.asm.tavg3d_mom_p.GEOS401.2002091500.2001091600.V01

6. File Collections

The contents of the Data Assimilation System file collections are described in sections 6.2, 6.3 and 6.4 while the sizes of each file are discussed in Section 6.1.

6.1 File Sizes

With the exception of chemistry (chm) products, GEOS-4 HDF files contain 1 day of data on a 1.25 degree x 1 degree lon-lat grid using 32-bit floating-point storage. Chm products are stored on a 2.5 degree by 2.0 degree lon-lat grid. See Appendix C for details about vertical levels. File sizes are approximate as metadata can cause slight variations in file size.

First-Look Filetype Group	Single Level (MB/day) (288,181,1) 8X/day	Pressure Levels (MB/day) (288,181,36) 4X/day	Eta Layers (MB/day) (288,181,55/24) 4X/day see appendix E
mis	31.7 (D4FAXMIS)	180.2 (D4FAPMIS)	551.6 (D4FAEMIS)
eng	51.7 (D4FAXENG)	N/A	N/A
str	21.7 (D4FAXSTR)	N/A	N/A
chm	0.4 (D4FAXCHM) *	8.8 (D4FAPCHM) **	N/A
cld	15.0 (D4FAXCLD)	Not produced	N/A
lsm	11.7 (D4FAXLSM)	N/A	N/A
mom	N/A	Not produced	N/A
mst	N/A	Not produced	N/A
tmp	N/A	Not produced	N/A
trp	N/A	Not produced	362.1 (D4FAETRP)
Totals	132.2	189.0	912.6

Late-Look Filetype Group	Single Level (MB/day) (288,181,1) 8X/day	Pressure Levels (MB/day) (288,181,36) 4X/day	Eta Layers (MB/day) (288,181,55/24) 4X/day see appendix E
mis	31.7 (D4LAXMIS)	180.2 (D4LAPMIS)	551.6 (D4LAEMIS)
eng	51.7 (D4LAXENG)	N/A	N/A
str	21.7 (D4LAXSTR)	N/A	N/A
chm	0.4 (D4LAXCHM) *	8.8 (D4LAPCHM) **	N/A
cld	15.0 (D4LAXCLD)	90.1 (D4LAPCLD)	N/A
lsm	11.7 (D4LAXLSM)	N/A	N/A
mom	N/A	150.2 (D4LAPMOM)	N/A
mst	N/A	60.1 (D4LAPMST)	N/A
tmp	N/A	150.2 (D4LAPTMP)	N/A
trp	N/A	120.1 (D4LAPTRP)	362.1 (D4LAETRP)
Total	132.2	759.7	912.6

* = (144, 91, 1) ** = (144, 91, 42)

6.2 Assimilated Synoptic Files

Below are the variables that are output into each syn file. These are instantaneous fields (no time averaging). The approximate size of each file below is determined by the following:

$$A \times B \times C \times D \times E \times F = \text{bytes/day}$$

where:

- A: X-Dimension
- B: Y-Dimension
- C: Vertical dimension
- D: Number of fields in file
- E: Number of times in file
- F: Number of bytes per floating point number

The method for calculating sizes is the same in 6.2, 6.3, and 6.4.

NOTE: All HDF variable names are UPPERCASE.

•**tsyn2d_mis_x** (8 times per day)

First-look ECS short name: D4FAXMIS

First-look ECS long name: DAS First-look 2d state (miscellaneous), instantaneous

Late-look ECS short name: D4LAXMIS

Late-look ECS long name: DAS Late-look 2d state (miscellaneous), instantaneous

Size: $288 \times 181 \times 1 \times 19 \times 8 \times 4 = 31.7 \text{ MB/day}$

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
PHIS	Surface geopotential	$(\text{m/s})^2$
PS	Surface pressure	hPa
SLP	Sea level pressure	hPa
SURFTYPE	Surface types	0=water, 1=land, 2=ice
TSKIN	Skin temperature	K
T2M	Temperature at 2 m above surface	K
T10M	Temperature at 10 m above surface	K
Q2M	Specific humidity at 2 m above surface	g/kg
Q10M	Specific humidity at 10 m above surface	g/kg
U2M	Zonal wind at 10 m above surface	m/s
V2M	Meridional wind at 2 m above surface	m/s
U10M	Zonal wind at 10 m above surface	m/s
V10M	Meridional wind at 10 m above surface	m/s
UFLUX	Zonal wind surface stress	N/m^2
VFLUX	Meridional wind surface stress	N/m^2
HFLUX	Sensible heat flux (pos. upward)	W/m^2
TROPP	Tropopause pressure	hPa
TROPT	Tropopause temperature	K
TROPQ	Tropopause specific humidity	g/kg

•**tsyn3d_mis_p** (4 times per day)

First-look ECS short name: D4FAPMIS

First-look ECS long name: DAS First-look 3d state (miscellaneous), instantaneous on pressure coordinates

Late-look ECS short name: D4LAPMIS

Late-look ECS long name: DAS Late-look 3d state (miscellaneous), instantaneous on pressure coordinates

Size: 288 x 181 x 36 x 6 x 4 x 4 = 180.2 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
UWND	Zonal wind	m/s
VWND	Meridional wind	m/s
HGHT	Geopotential height (virtual)	m
TMPU	Temperature	K
SPHU	Specific humidity	g/kg
RH	Relative humidity	percent

6.3 Assimilated Time Averaged Files

Below are the variables that are output in each "tavg" file. These are time-averaged fields. Single-level, or 2-dimensional data will be output every 3 hours while 3-dimensional data will be output every 6 hours.

•**tavg2d_eng_x** (8 times per day)

First-look ECS short name: D4FAXENG

First-look ECS long name: DAS First-look 2d energy, averaged

Late-look ECS short name: D4LAXENG

Late-look ECS long name: DAS Late-look 2d energy, averaged

Size: 288 x 181 x 1 x 31 x 8 x 4 = 51.7 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
PREACC	Total precipitation rate	mm/day
PRECL	Large-scale precipitation rate	mm/day
PRECON	Convective precipitation rate	mm/day
TPW	Total precipitable water	g/cm ²
EVAP	Surface evaporation	mm/day
GWETROOT	Root zone soil wetness	fraction
GWETTOP	Top soil layer wetness	fraction
HFLUX	Sensible heat flux (positive upward)	W/m ²
TSKIN	Skin temperature	K
T2M	Temperature interpolated to 2 meters	K
T10M	Temperature interpolated to 10 meters	K
Q2M	Specific humidity interpolated to 2 meters	g/kg
Q10M	Specific humidity interpolated to 10 meters	g/kg
RADLWG	Net upward longwave flux at the ground	W/m ²
RADSWG	Net downward shortwave flux at the ground	W/m ²
PARDF	Diffuse-beam photosynthetically active radiation	W/m ²
PARDR	Direct-beam photosynthetically active radiation	W/m ²
LWGCLR	Clear sky net longwave flux at the ground	W/m ²
SWGCLR	Clear sky net downward shortwave radiation at the ground	W/m ²
ALBEDO	Surface albedo	fraction
ALBVISDR	Direct beam VIS surface albedo	fraction
ALBVISDF	Diffuse beam VIS surface albedo	fraction
ALBNIRDR	Direct beam NIR surface albedo	fraction
ALBNIRDF	Diffuse beam NIR surface albedo	fraction
VAVEU	Vertically averaged zonal wind	m/s
VAVEV	Vertically averaged meridional wind	m/s
VAVET	Vertically averaged temperature	K
VAVEUQ	Vertically averaged UWND*SPHU	m-g/s-kg
VAVEVQ	Vertically averaged VWND*SPHU	m-g/s-kg

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
VAVEUT	Vertically averaged UWND*TMPU	m-K/s
VAVEVT	Vertically averaged VWND*TMPU	m-K/s

•**avg2d_str_x** (8 times per day)

First-look ESDT: D4FAXSTR

First-look ECS long name: DAS First-look 2d stress, averaged

Late-look ECS short name: D4LAXSTR

Late-look ECS long name: DAS Late-look 2d stress, averaged

Size: 288 x 181 x 1 x 13 x 8 x 4 = 21.7 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
PS	Time averaged surface pressure	hPa
UFLUX	Zonal wind surface stress	N/m ²
VFLUX	Meridional wind surface stress	N/m ²
GWDUS	Zonal wind gravity wave surface stress	N/m ²
GWDVS	Meridional wind gravity wave surface stress	N/m ²
USTAR	Friction velocity	m/s
Z0H	Roughness length, sensible heat	m
Z0M	Roughness length, momentum	m
PBLH	Planetary boundary layer height	m
U2M	Zonal wind interpolated to 2 meters	m/s
V2M	Meridional wind interpolated to 2 meters	m/s
U10M	Zonal wind interpolated to 10 meters	m/s
V10M	Meridional wind interpolated to 10 meters	m/s

•**avg2d_lsm_x** (8 times per day)

First-look ESDT: D4FAXLSM

First-look ECS long name: DAS First-look 2d Land Surface Model (LSM), averaged

Late-look ECS short name: D4LAXLSM

Late-look ECS long name: DAS Late-look 2d Land Surface Model (LSM), averaged

Size: 288 x 181 x 1 x 7 x 8 x 4 = 11.7 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
RAINCON	Convective rainfall (liquid precipitate)	mm/day
RAINLSP	Large-scale rainfall (liquid precipitate)	mm/day
LWGDOWN	Downward longwave radiation at the ground	W/m ²
LWGUP	Upward longwave radiation at the ground	W/m ²
LAI	Leaf area index	percent
DTG	Total change in ground temperature	K/s
SNOW	Snow depth	mm (water equivalent)

•**avg2d_cld_x** (8 times per day)

First-look ESDT: D4FAXCLD
 First-look ECS long name: DAS First-look 2d cloud, averaged
 Late-look ECS short name: D4LAXCLD
 Late-look ECS long name: DAS Late-look 2d cloud, averaged
 Size: 288 x 181 x 1 x 9 x 8 x 4 = 15.0 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
OLR	Outgoing longwave radiation	W/m ²
OLRCLR	Clear sky outgoing longwave radiation	W/m ²
RADSWT	Incident shortwave radiation at top of atmosphere	W/m ²
OSR	Outgoing shortwave radiation	W/m ²
OSRCLR	Clear sky outgoing shortwave radiation	W/m ²
CLDFRC	Total cloud fraction	fraction
CLDLW	Low-level (1000-700 hPa) cloud fraction	fraction
CLDMID	Mid-level (700-400 hPa) cloud fraction	fraction
CLDHI	High-level (above 400 mb) cloud fraction	fraction

•**avg3d_cld_p** (4 times per day)

Late-look ECS short name: D4LAPCLD
 Late-look ECS long name: DAS Late-look 3d cloud, averaged on pressure coordinates
 Size: 288 x 181 x 36 x 3 x 4 x 4 = 90.1 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
TAUCLI	Cloud optical depth, ice	non-dimensional
TAUCLW	Cloud optical depth, water	non-dimensional
CLDTOT	3-D total cloud fraction	fraction

•**avg3d_mom_p** (4 times per day)

Late-look ECS short name: D4LAPMOM
 Late-look ECS long name: DAS Late-look 3d momentum, averaged on pressure coordinates
 Size: 288 x 181 x 36 x 5 x 4 x 4 = 150.2 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
TURBU	Zonal wind tendency due to turbulence	m/s/day
TURBV	Meridional wind tendency due to turbulence	m/s/day
GWDU	Zonal wind tendency due to gravity wave drag	m/s/day
GWDV	Meridional wind tendency due to gravity wave drag	m/s/day
OMEGA	Vertical velocity	Pa/s

•**tavg3d_mst_p** (4 times per day)

Late-look ECS short name: D4LAPMST

Late-look ECS long name: DAS Late-look 3d moisture, averaged on pressure coordinates

Size: $288 \times 181 \times 36 \times 2 \times 4 \times 4 = 60.1$ MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
TURBQ	Specific humidity tendency due to turbulence	g/kg/day
MOISTQ	Specific humidity tendency due to moist processes	g/kg/day

•**tavg3d_tmp_p** (4 times per day)

Late-look ECS short name: D4LAPTMP

Late-look ECS long name: DAS Late-look 3d temperature, averaged on pressure coordinates

Size: $288 \times 181 \times 36 \times 5 \times 4 \times 4 = 150.2$ MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
TURBT	Temperature tendency due to turbulence	K/day
MOISTT	Temperature tendency due to moist processes	K/day
RADLW	Temperature tendency due to longwave radiation	K/day
RADSW	Temperature tendency due to shortwave radiation	K/day
GWDT	Temperature tendency due to gravity wave drag	K/day

•**tavg3d_trp_p** (nominally 4 times per day)

Late-look ECS short name: D4LAPTRP

Late-look ECS long name: DAS Late-look 3d transport, averaged on pressure coordinates

Size: $288 \times 181 \times 36 \times 4 \times 4 \times 4 = 120.1$ MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
KH	Eddy diffusivity coefficient for scalars	m^2/s
KM	Eddy diffusivity coefficient for momentum	m^2/s
CLDMAS	Cloud mass flux	$\text{kg}/\text{m}^2/\text{s}$
DTRAIN	Detrainment cloud mass flux	$\text{kg}/\text{m}^2/\text{s}$

6.4 Assimilated Chemistry Files

The following variables are generated by the GMAO Ozone Assimilation System. These are instantaneous fields (no time averaging). These products are produced from an offline (OL) ozone assimilation, which uses the wind data from the DAS as input. The ozone assimilation operates at a coarser resolution than the DAS, thus these products have horizontal resolution of 2.5 x 2 degrees. In addition, the vertical structure of the 3-D ozone products is different than other 3-D DAS products. See Appendix C for details.

The ozone mixing ratio fields in the tavg3d_chm_p product at the following levels: 1000, 975, 950, 925, 900, and 875 hPa are provided at the request of users. The ozone values provided at these levels are identical to the values at 850 hPa. Ozone data are assimilated between 850 and 0.2 hPa. No geophysical significance should be attributed to the 1000, 975, 950, 915, 900, and the 875 hPa levels.

•tsyn2d_chm_x (8 times per day)

First-look ECS short name: D4FAXCHM

First-look ECS long name: DAS First-look 2d chemistry, instantaneous

Late-look ECS short name: D4LAXCHM

Late-look ECS long name: DAS Late-look 2d chemistry, instantaneous

Size: 144 x 91 x 1 x 1 x 8 x 4 = 0.4 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
OZONE	Total ozone	DU

•tsyn3d_chm_p (4 times per day)

First-look ECS short name: D4FAPCHM

First-look ECS long name: DAS First-look 3d chemistry, instantaneous on pressure coordinates

Late-look ECS short name: D4LAPCHM

Late-look ECS long name: DAS Late-look 3d chemistry, instantaneous on pressure coordinates

Size: 144 x 91 x 42 x 1 x 4 x 4 = 8.8 MB/day

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
OZONE	Ozone mixing ratio	ppmv

7. Metadata

GEOS-4 gridded output files will be associated with two types of metadata. Depending on the utility you use to access the file, one set of metadata will be read and the other ignored.

7.1 EOSDIS Metadata

If you are using the EOSDIS toolkit you will only use the EOSDIS metadata. EOSDIS identifies two major types of metadata, collection and granule.

Collection metadata are stored in a separate index file. This file is like a library card catalog. Each ESDT has a "card" that contains its unique collection attributes. Appendix B describes the ESDT collection metadata.

Granule metadata is the "table of contents" information stored on the data file itself. The EOSDIS granule metadata include:

- file name (local granule ID)
- grid structure
- number of times per day fields are stored in this file
- number of vertical levels for each variable in this file
- names of variables in this file
- variable format (32-bit floating point, 16-bit integer, etc.)
- variable storage dimensions
 - 2-d fields will have 3 storage dimensions, time, latitude and longitude
 - 3-d fields will have 4 storage dimensions, time, latitude, longitude and vertical levels
- "missing" value for each variable
- unpacking scale factor for each packed variable (see section 8)
- unpacking off-set value for each packed variable (see section 8)

7.2 COARDS Metadata

If you use GRADS or FERRET to view GEOS-4 gridded data sets you will only use the COARDS metadata. These metadata will comply with the COARDS convention and include the following information:

- space-time grid information (dimension variables)
- variable long names (descriptions)

- variable units
- "missing" value for each variable
- unpacking scale factor for each packed variable (see section 8)
- unpacking off-set value for each packed variable (see section 8)

8. Sample Software

Presented here is software that illustrates using the standard HDF library or the ECS HDF-EOS library to read GEOS-4 products. The program shown below will accept as command line arguments a file name and a field name. It will open the file, read the requested field at the first time, compute an average for this field, and print the result to standard output. There are two versions of this program. The first version uses the HDF-EOS library to read the file. The second version uses the standard HDF library to read the file. Electronic copies of these programs can be obtained from the Operations section of the GMAO web page:

<http://gmao.gsfc.nasa.gov/Operations/>

```

/*****
/* This program demonstrates how to read a field from a GMAO HDF-EOS */
/* product using the HDF-EOS library. It will take a file name and */
/* field name on the command line, read the first time of the given */
/* field, calculate an average of that time and print the average. */
/* */
/* usage: avg <file name> <field name> */
/* */
/* Rob Lucchesi */
/* rlucchesi@GMAO.gsfc.nasa.gov */
/* 2/12/1999 */
*****/

#include "hdf.h"
#include "mfhdf.h"
#include <stdio.h>

#define XDIM 288
#define YDIM 181
#define ZDIM 36

main(int argc, char *argv[]) {
int32 sd_id, sds_id, status;
int32 sds_index;
int32 start[4], edges[4], stride[4];
char *fname, *vname;
float32 data_array[ZDIM][YDIM][XDIM];
float32 avg, sum;
int32 i,j,k;
int32 file_id, gd_id;

if (argc != 3) {

```

```
printf("Usage: avg <filename> <field> \n");
exit (-1);
}

fname = argv[1];
vname = argv[2];

/* Open the file (read-only) */

file_id = GDopen (fname, DFACC_RDONLY);
if (file_id < 0) {
printf ("Could not open %s\n",fname);
exit(-1);
}

/* Attach to the EOS grid contained within the file. */
/* The GMAO uses the generic name "EOSGRID" for the grid in all products. */

gd_id = GDattach (file_id,"EOSGRID");
if (gd_id < 0) {
printf ("Could not open %s\n",fname);
exit(-1);
}

/* Set positioning arrays to read the entire field at the first time. */

start[0] = 0;
start[1] = 0;
start[2] = 0;
start[3] = 0;

stride[0] = 1;
stride[1] = 1;
stride[2] = 1;
stride[3] = 1;

edges[0] = 1;
edges[1] = ZDIM;
edges[2] = YDIM;
edges[3] = XDIM;

/*
```

In this program, we read the entire field. By manipulating the start and edges arrays, it is possible to read a subset of the entire array. For example, to read a 3D section defined by $x=100,224$; $y=50,149$; $z=15,16$ you would set the start and edges arrays to the following:

```
start[0] = 0; time start location
start[1] = 15; z-dim start location
start[2] = 50; y-dim start location
start[3] = 100; x-dim start location
```

```
edges[0] = 1; time length
edges[1] = 2; z-dim length
edges[2] = 100; y-dim length
edges[3] = 125; x-dim length
```

```
*/
```

```
/* Read the data into data_array */
```

```
status = GDreadfield (gd_id, vname, start, stride, edges, data_array);
printf ("Read status=%d\n",status);
```

```
/* Calculate and print the average */
```

```
sum=0.0;
for (i=0; i<XDIM; i++)
for (j=0; j<YDIM; j++)
for (k=0; k<ZDIM; k++)
sum += data_array[k][j][i];
```

```
avg = sum/(XDIM*YDIM*ZDIM);
```

```
printf ("Average of %s in 3 dimensions is=%f\n",vname,avg);
```

```
/* Close file. */
```

```
status = GDdetach (gd_id);
status = GDclose (file_id);
```

```
}
```

```

/*****
/* This program demonstrates how to read a field from a GMAO HDF-EOS */
/* product using the HDF library (HDF-EOS not required). It will take */
/* a file name and field name on the command line, read the first time */
/* of the given field, calculate an average of that time and print the average. */
/* */
/* usage: avg <file name> <field name> */
/* */
/* Rob Lucchesi */
/* rlucchesi@gmao.gsfc.nasa.gov */
/* 2/12/1999 */
*****/

#include "hdf.h"
#include "mfhdf.h"
#include <stdio.h>

#define XDIM 288
#define YDIM 181
#define ZDIM 36

main(int argc, char *argv[]) {
int32 sd_id, sds_id, status;
int32 sds_index;
int32 start[4], edges[4], stride[4];
char *fname, *vname;
float32 data_array[ZDIM][YDIM][XDIM];
float32 avg, sum;
int32 i,j,k;

if (argc != 3) {
printf("Usage: avg <filename> <field> \n");
exit (-1);
}

fname = argv[1];
vname = argv[2];

/* Open the file (read-only) */

sd_id = SDstart (fname, DFACC_RDONLY);
if (sd_id < 0) {
printf ("Could not open %s\n", fname);

```

```
exit(-1);
}
```

```
/* Find the index and ID of the SDS for the given variable name. */
```

```
sds_index = SDnametoindex (sd_id, vname);
if (sds_index < 0) {
printf ("Could not find %s\n",vname);
exit(-1);
}
```

```
sds_id = SDselect (sd_id,sds_index);
```

```
/* Set positioning arrays to read the entire field at the first time. */
```

```
start[0] = 0;
start[1] = 0;
start[2] = 0;
start[3] = 0;
```

```
stride[0] = 1;
stride[1] = 1;
stride[2] = 1;
stride[3] = 1;
```

```
edges[0] = 1;
edges[1] = ZDIM;
edges[2] = YDIM;
edges[3] = XDIM;
```

```
/*
```

In this program, we read the entire field. By manipulating the start and edges arrays, it is possible to read a subset of the entire array. For example, to read a 3D section defined by x=100,224; y=50,149; z=15,16 you would set the start and edges arrays to the following:

```
start[0] = 0; time start location
start[1] = 15; z-dim start location
start[2] = 50; y-dim start location
start[3] = 100; x-dim start location
```

```
edges[0] = 1; time length
edges[1] = 2; z-dim length
```

```
edges[2] = 100; y-dim length
edges[3] = 125; x-dim length

*/

/* Read the data into data_array */

status = SDreaddata (sds_id, start, stride, edges, (VOIDP) data_array);
printf ("read status=%d\n",status);

/* Calculate and print the average */

sum=0.0;
for (i=0; i<XDIM; i++)
for (j=0; j<YDIM; j++)
for (k=0; k<ZDIM; k++)
sum += data_array[k][j][i];

avg = sum/(XDIM*YDIM*ZDIM);

printf ("Average of %s in 3 dimensions is=%f\n",vname,avg);

/* Close file. */

status = SDendaccess (sds_id);
status = SDend (sd_id);

}
```

Appendix A. Types of Assimilation Configurations

First Look Assimilation. Atmospheric observations from satellites, balloons, aircraft, ships, and other sources are grouped into six-hour data windows and processed by the atmospheric analysis four times each day. The first look analysis will run 6 to 10 hours after the 4 analysis times (0Z, 6Z, 12Z, 18Z). It will run using whatever conventional and satellite observations are available at the data cut-off time. Even though first-look will run as 4 distinct segments, the GMAO will only distribute products that contain one complete day. Products produced from this and any other *assimilation* are a combination of output from the statistical analysis system and a short GCM forecast.

Late Look Assimilation. The software configuration is identical to the first look assimilation but, due to a delay of at least two weeks, a more complete set of input observations can be used. Unlike the first look analysis, the late look analysis will process one day at a time. The late look assimilation produces a greater volume of output products, since computational performance is not as critical.

Forecast/Simulation. This is a GCM forecast, with no insertion of atmospheric data via the analysis. The only outside data that enters the system are the boundary conditions, i.e., sea surface temperature and sea-ice concentration. Five-day forecasts are typically generated to support NASA field campaigns and to assess assimilation and forecast skill. Multi-year simulations are produced to investigate the climatology of the GCM. GMAO forecast products are not distributed to ECS and file formats are not discussed in this document.

Reprocessing. The GMAO will reprocess specified time periods since EOS-Terra launch using a recent version of the GEOS DAS software to support instrument team reprocessing requirements. A reprocessing run will use the Late Look assimilation file groupings and will use a version of GEOS DAS that is held constant through the reprocessing period. It is expected that new ECS ESDTs will be generated for each reprocessing run.

Reanalysis. The GMAO will occasionally run reanalysis experiments. Reanalysis is the same as reprocessing except the time period is often much longer and not necessarily part of the EOS period. Reanalysis experiments are often run using baseline versions of the GEOS DAS system to support a wide variety of research activities internal and external to the GMAO. Unique ESDTs will be generated for any reanalysis data distributed through ECS.

Appendix B. Collection Metadata

GEOS-4 collection metadata will contain the following. To view the ESDTs associated with GMAO products, which include collection metadata, see the GMAO Operations web page:

<http://gmao.gsfc.nasa.gov/Operations>.

ECS Collection

- Revision Date
- Suggested Usage

Single Type Collection

- Collection State
- Maintenance and Update Frequency

Spatial

- Spatial Coverage Type

Bounding Rectangle

- West Bounding Coordinate
- North Bounding Coordinate
- East Bounding Coordinate
- South Bounding Coordinate

Altitude System Definition (for 3d files only)

- Altitude Datum Name
- Altitude Distance Units
- Altitude Encoding Method
- Altitude Resolution Class
- Altitude Resolution

Depth System Definition (land surface files only)

- Depth Datum Name
- Depth Distance Units
- Depth Encoding Method
- Depth Resolution Class
- Depth Resolution

Geographic Coordinate System

- Latitude Resolution
- Longitude Resolution
- Geographic Coordinate Units

Temporal

- Time Type

Date Type
Temporal Range Type
Precision of Seconds
Ends at Present Flag

Range Date Time

Range Beginning Date
Range Beginning Time
Range Ending Date
Range Ending Time

Contact Person

Role
Hours of Service
Contact Job Position
Contact First Name
Contact Middle Name
Contact Last Name

Contact Person Address

Street address
City
State/Province
Postal Code
Country

Telephone

Telephone Container
Telephone Number
Telephone Number Type

Email

Electronic Mail Address

Contact Organization

Role
Hours of Service
Contact Instruction
Contact Organization Name

Contact Organization Address

Street Address
City

State/Province
Postal Code
Country

Organization Telephone Number
Telephone Number
Telephone Number Type

Organizational Email
Electronic Mail Address

Discipline Topic Parameters
ECS Discipline Keyword
ECS Topic Keyword
ECS Term Keyword
ECS Variable Keyword
ECS Parameter Keyword

Temporal Keyword Class
Temporal Keyword

Spatial Keyword Class
Spatial Keyword

Processing Level
Processing Level Description
Processing Level ID

Analysis Source
Analysis Short Name
Analysis Long Name
Analysis Technique
Analysis Type

CSDT Description
Primary CSDT

Additional Attributes
Additional Attribute Data Type
Additional Attribute Description
Additional Attribute Name

Physical Parameter Details

- Parameter Units of Measure
- Parameter Range
- Parameter Value Accuracy
- Parameter Value Accuracy Explanation
- Parameter Measurement Resolution

Storage Medium Class (filled in by DAAC)

- Storage Medium

Appendix C. Vertical Grid Structure

Pressure level data will be output on the following 36 pressure levels for all 3-D products except chemistry:

Level	Pressure (hPa)	Level	Pressure (hPa)	Level	Pressure (hPa)
1	1000	13	600	25	50
2	975	14	550	26	40
3	950	15	500	27	30
4	925	16	450	28	20
5	900	17	400	29	10
6	875	18	350	30	7
7	850	19	300	31	5
8	825	20	250	32	3
9	800	21	200	33	2
10	750	22	150	34	1
11	700	23	100	35	0.4
12	650	24	70	36	0.2

Due to the scientific characteristics of the ozone assimilation, 3-D chemistry products (D4FAPCHM, D4LAPCHM) will have a different set of pressure levels:

Level	Pressure (hPa)	Level	Pressure (hPa)	Level	Pressure (hPa)
1	1000	15	500	29	50
2	975	16	450	30	40
3	950	17	400	31	30
4	925	18	350	32	25
5	900	19	300	33	20
6	875	20	250	34	15
7	850	21	200	35	10
8	825	22	170	36	7
9	800	23	150	37	5
10	750	24	130	38	3
11	700	25	115	39	2
12	650	26	100	40	1
13	600	27	85	41	0.4
14	550	28	70	42	0.2

Appendix D. Summary of differences between GEOS-3 and GEOS-4

In this section we summarize the differences between the GEOS-3 and GEOS-4 products. We will not discuss the scientific characteristics of the systems, but only the differences in format and content of the products. GEOS-3 products are generated using the GEOS-3 Data Assimilation System until October 1, 2002, after which products from the GEOS-4 Data Assimilation System will be produced. The same 21 products are produced from both systems, 8 from First-look and 13 from Late-look. The format of all products is the HDF-4 version of HDF-EOS. Details of the differences follow.

- The horizontal resolution is different. The GEOS-3 resolution is 360x181. The GEOS-4 resolution is 288x181.
- The averaging periods of time-averaged products has been changed. Averages are now between synoptic times rather than spanning them. For example, the first average in a 3D 6-hourly file from GEOS-4 is the period from 00 GMT to 06 GMT. The first average in a corresponding GEOS-3 file would be 21 GMT to 03 GMT. This has changed the effective range times of the GEOS-4 time-averaged products. See section 3 of this document for more information.
- The contents of most products changed.

○tsyn2d_mis_x:

- ALBEDO and TPW removed but still available as time-averaged in tavg2d_eng_x
- VAVEU, VAVEV, and VAVET moved to tavg2d_eng_x, now time-averaged
- SNOW removed but still available as time-averaged in tavg2d_lsm_x
- GWET moved to tavg2d_eng_x as GWETTOP and GWETROOT, now time-averaged
- TGROUND renamed to TSKIN
- TROPQ added

otsyn3d_mis_p:

- TKE removed
- OMEGA moved to tavg3d_mom_p, now time-averaged

otavg2d_eng_x:

- QICE, CT, VAVEQIAU, VAVEQFIL, and VAVETIAU removed
- VAVEU, VAVEV, VAVET, GWETTOP, and GWETROOT added (formerly in tsyn2d_mis_x)
- TGROUND renamed to TSKIN
- PRECL added
- PARDR and PARDF added (formerly in tavg2d_lsm_x)

otavg2d_str_x:

- GWDUT, GWDVT, CU, and PIAU removed
- Z0 split into Z0H and Z0M
- PBL changed to PBLH (units changed from hPa to m)

otavg2d_lsm_x:

- SNOWFALL, GREEN, and DLWDTC removed
- PARDR and PARDF moved to tavg2d_eng_x

otavg2d_cld_x:

- TAULOW, TAUMID, TAUHI, CLDTMP, and CLDPRS removed

otavg3d_cld_p:

- CLDRAS removed

- TAUCLD split into TAUCLI and TAUCLW
- **tavg3d_mom_p**:
 - RFU, RFV, UIAU, VIAU, DUDT, and DVDT removed
 - OMEGA was added (formerly in tsyn3d_mis_p)
- **tavg3d_mst_p**:
 - DQLS, QIAU, QFILL, and DQDT removed
- **tavg3d_tmp_p**:
 - TURBT, MOISTT, DTLS, RADLW, RADSW, RFT, GWDT, TIAU, DTDI, LWCLR, and SWCLR removed
- Due to these changes, there are differences in the daily file sizes.
- The filename convention was changed slightly. *AMI* was changed to *GEOS4*. The hour in the filename is different for time-averaged products due to the difference of time ranges mentioned above. See section 5 for more details about the file naming convention.
- New ESDTs were defined for GEOS-4. The names are listed in section 6 of this document. For First-look the ESDT names now start with *D4F* and for Late-look *D4L*. For GEOS-3 the names started with *DFL* and *DLL*.
- The data contained in the tsyn2d_mis_x product is 3-hourly, however the source data from GOES-4 is 6-hourly. The interim times are computed by time interpolating between synoptic times. This is done to provide consistency with the corresponding GEOS-3 product. See section 3 for more information.

Appendix E: Additional products on eta vertical coordinates

A set of First-look (FLK) and Late-look (LLK) products defined on the GEOS-4 vertical coordinate grid (eta) will be produced starting in spring 2004. Eta coordinates are generically known as atmosphere hybrid sigma pressure coordinates. The two eta-layer products will be daily files with four 6-hour time-averages for each variable. These files are targeted for users involved in chemical transport work, but may be useful to a wider audience. The *tavg3d_mis_e* product is defined for 55 vertical layers, from the highest layer of the GCM (0.015 hPa) to the layer adjacent to the surface. The *tavg3d_trp_e* product is defined for only the lowest 24 GCM eta layers because the variables included vanish above the tropopause. The time period for each six-hour average will be centered on one of the 4 synoptic times, i.e., 00 GMT will represent the average from 21 GMT to 03 GMT. This is different than the pressure products, described in section 6, which span the periods between synoptic times. For more information, please see the diagram at the end of this appendix.

The GEOS-4 eta coordinate system can be described by the following equation:

$$p(i,j,k,t) = a(k) + b(k)*ps(i,j,t)$$

where for a given time t , p is the pressure at the coordinates i,j,k and $a(k)$ and $b(k)$ are components of the hybrid coordinate at the layer edge k . To find the average p for a layer, one would average p at the edges k and $k+1$. $ps(i,j,t)$ is the surface pressure at the horizontal gridpoint (i,j) and time t . $a(k)$ and $b(k)$ at 56 layer edges (25 edges for the 24 layer product) are stored as global metadata values in each eta product file. They are also listed in a table near the end of this appendix. For *tavg3d_mis_e*, when $k=1$, $p = 1.0$ Pa and when $k=56$, $p=ps$. For *tavg3d_trp_e*, when $k=1$, $p=3,381$ Pa and when $k=56$, $p=ps$.

•**tavg3d_mis_e** (4 times per day)

FLK ECS short name: D4FAEMIS

FLK ECS long name: DAS First-look miscellaneous meteorology fields, time-averaged on eta coordinates

LLK ECS short name: D4LAEMIS

LLK ECS long name: DAS Late-look miscellaneous meteorology fields, time-averaged on eta coordinates

Size: 551 MB/day

Dimensions:

longitude: 288

latitude: 181

vertical layers (eta): 55

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
PS*	Surface pressure (two-dimensional field)	Pa
DIABDT	T Tendency - Total diabatic (physics)	K/s
HGHT	Geopotential height at mid-layer	m
HGHTTE	Geopotential height at layer top	m

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
KH	Vertical diffusion diffusivities (heat/moisture)	m ² /s
KM	Eddy diffusivity for momentum	m ² /s
OMEGA	Vertical pressure velocity	Pa/s
PV	Ertels potential vorticity	K*m ² /(kg*sec)
RH	Relative humidity after cloud physics	percent
SPHU*	Specific humidity	kg/kg
TMPU	Temperature	K
UWND	U wind	m/s
VWND	V wind	m/s

•**tavg3d_trp_e** (4 times per day)

FLK ECS short name: D4FAETRP

FLK ECS long name: DAS First-look transport fields, time-averaged on eta coordinates

LLK ECS short name: D4LAETRP

LLK ECS long name: DAS Late-look transport fields, time-averaged on eta coordinates

Size: 382 MB/day

Dimensions:

longitude: 288

latitude: 181

vertical layers (eta): 24

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
PS*	Surface pressure (two-dimensional field)	Pa
CLDTOT	3-D Cloud fraction	fraction
CMFDQR2	Rain production rate - Hack convection	kg/kg/s
CMFDTR	Detrainment mass flux - Hack convection	Pa/s
CMFETR	Entrainment mass flux - Hack convection	Pa/s
CMFMC	Total moist convection mass flux	Pa/s
CMFMC2	Hack moist convection mass flux	Pa/s
DQRL	Rain production rate - large-scale	kg/kg/s
DTRAIN*	Detrainment cloud mass flux (Ave)	Pa/s
HKBETA	Overshoot parameter in Hack scheme	fraction
HKETA	Mass flux without overshoot in Hack scheme	kg/m ² /s
MOISTQ	Q tendency - moist physics	kg/kg/s
TAUCLI	Cloud optical depth ice	non-dimensional
TAUCLW	Cloud optical depth water	non-dimensional
ZMDQR	Rain production rate - Z&M convection	kg/kg/s
ZMDU	Updraft detrainment mass flux - Z&M convection	Pa/s
ZMED	Downdraft entrainment mass flux - Z&M convection	Pa/s
ZMEU	Updraft entrainment mass flux - Z&M convection	Pa/s

<u>Variable Name</u>	<u>Description</u>	<u>Units</u>
ZMMD	Downdraft mass flux - Z&M convection (Ave)	Pa/s
ZMMU	Updraft mass flux - Z&M convection	Pa/s

* - These variables are also available in other GEOS-4 surface or pressure products, but in the eta products we use standard SI units or derivations. In particular, tendency terms in pressure level products are expressed as rates of change per day, while the eta products are per second. Moisture concentrations are expressed here as kg/kg, while the pressure products use g/kg. Surface pressure is expressed in Pa instead of hPa. Mass flux terms are expressed as Pa/s.

Filenames of the eta products conform to the conventions described in section 5 of this document. The beginning and ending times will be slightly different than for the pressure products due to the differences in the time-averaging period. Here is an example of filenames from First-look.

DAS.flk.asm.tavg3d_mis_e.GEOS403.2003070121.2003070221.V01
 DAS.flk.asm.tavg3d_trp_e.GEOS403.2003070121.2003070221.V01

The following table contains ak and bk pairs for each layer edge k. There are 56 layer edges that define 55 layers. Note that for the 24 layer *tavg3d_trp_e* product, there are 25 layer edges and the k values are denoted with ()s. These arrays can be found as floating point global metadata attributes in each eta product file.

k	ak	bk		k	ak	bk
1	1.000	0.000		29	1979.155	0.000
2	2.000	0.000		30	2373.036	0.000
3	3.270	0.000		31	2836.782	0.000
4	4.759	0.000		32 (1)	3380.996	0.000
5	6.600	0.000		33 (2)	4017.542	0.000
6	8.934	0.000		34 (3)	4764.393	0.000
7	11.970	0.000		35 (4)	5638.794	0.000
8	15.950	0.000		36 (5)	6660.338	0.000
9	21.135	0.000		37 (6)	7851.230	0.000
10	27.853	0.000		38 (7)	9236.566	0.000
11	36.504	0.000		39 (8)	10866.343	0.000
12	47.581	0.000		40 (9)	12783.700	0.000
13	61.678	0.000		41 (10)	15039.300	0.000
14	79.513	0.000		42 (11)	17693.000	0.000
15	101.944	0.000		43 (12)	20119.209	0.007
16	130.051	0.000		44 (13)	21686.492	0.028
17	165.079	0.000		45 (14)	22436.287	0.064

k	ak	bk		k	ak	bk
18	208.497	0.000		46 (15)	22388.469	0.115
19	262.021	0.000		47 (16)	21541.752	0.183
20	327.643	0.000		48 (17)	19873.783	0.270
21	407.657	0.000		49 (18)	17340.318	0.378
22	504.680	0.000		50 (19)	13874.440	0.510
23	621.680	0.000		51 (20)	10167.165	0.643
24	761.984	0.000		52 (21)	6609.843	0.765
25	929.294	0.000		53 (22)	3546.596	0.868
26	1127.689	0.000		54 (23)	1270.494	0.943
27	1364.339	0.000		55 (24)	0.000	0.985
28	1645.707	0.000		56 (25)	0.000	1.000




Figure 1 - This is a graphical representation of the averaging periods for GEOS-4 3D time-averaged products. Note that eta-layer products are centered on synoptic times while pressure-level products span the range between synoptic times.

REFERENCES

GMAO Operations web page: <http://gmao.gsfc.nasa.gov/Operations/>

DAO, 2002: Algorithm Theoretical Basis Document, Data Assimilation Office, NASA's Goddard Space Flight Center.

Doplick, T., 1997: The Role of Metadata in EOSDIS (DRAFT). EOSDIS Technical Paper 160-TP-013-001.

Gross, C., 1997: B.0 Earth Science Data Model for the ECS Project. EOSDIS Technical Paper 420-TP-015-001.

NOAA, 1995: Conventions for the Standardization of NetCDF Files (COARDS). http://ferret.wrc.noaa.gov/noaa_coop/coop_cdf_profile.html

NCSA, 1999: The NCSA HDF Home Page. <http://hdf.ncsa.uiuc.edu/>

NASA, 1999: HDF-EOS Standards. <http://hdfeos.gsfc.nasa.gov/>

deWitt, Shaun, 1996: Writing HDF-EOS Grid Products for Optimum Subsetting Services. ECS Technical Paper 170-TP-007-001.